ISSN (print) 0093-4666 © 2015. Mycotaxon, Ltd. ISSN (online) 2154-8889

MYCOTAXON

http://dx.doi.org/10.5248/130.421

Volume 130, pp. 421–427 *April–June 2015*

Molecular analyses reveal a new species in *Melanoderma* from tropical China

Hai-Sheng Yuan [1*] & Yu-He Kan [1, 2]

[1] *State Key Laboratory of Forest and Soil Ecology, Institute of Applied Ecology, Chinese Academy of Sciences, Shenyang 110164, P. R. China*

[2] *Graduate University of the Chinese Academy of Sciences, Beijing 100049, P. R. China*

* Correspondence to: *hsyuan@iae.ac.cn*

Abstract — A new wood-decaying polypore, *Melanoderma disciforme*, is described from tropical China based on morphological characteristics and rDNA ITS sequences. The new species is characterized by perennial effuse-reflexed basidiocarps, a narrow pileus with a thin black crust on the pileal surface, a dimitic hyphal system with dextrinoid and cyanophilous skeletal hyphae, and the presence of fusiform cystidioles and cylindrical basidiospores. Molecular analyses indicate that the new species and *M. microcarpum* cluster together with full support.

Key words —*Basidiomycota*, *Polyporaceae*, phylogeny, taxonomy

Introduction

The genus *Melanoderma* B.K. Cui & Y.C. Dai was recently established to accommodate the newly described species *M. microcarpum* B.K. Cui & Y.C. Dai (Cui et al. 2011). The genus is characterized by perennial pileate to effuse-reflexed basidiocarps, a black pileal surface with a thin crust, a dimitic hyphal system with dextrinoid cyanophilous skeletal hyphae, apically encrusted cystidia, and the presence of cystidioles and cylindrical thin-walled basidiospores. Microscopically, *Melanoderma* resembles *Perenniporia* Murrill in having dextrinoid and cyanophilous skeletal hyphae. However, *Melanoderma* is distinguished from *Perenniporia* by its distinctly thin-walled, non-dextrinoid, and acyanophilous basidiospores (Cui et al. 2011).

China is very rich in wood-inhabiting fungi species, and recently extensive studies have been conducted on species diversity, taxonomy, ecological patterns, and phylogeny of wood-inhabiting fungi (Dai 2010, 2011, 2012, Dai et al. 2007, Ma et al. 2011, Zhang et al. 2009, Zhou & Dai 2012). Two polypore specimens collected in 2005 from a tropical forest of Yunnan Province were

subsequently identified and deposited as *Melanoderma microcarpum*. But recent comprehensive morphological and molecular analyses suggest that these specimens represent an undescribed *Melanoderma* species. Here we clarify its taxonomic status and describe it as a new species.

Materials & methods

Morphological studies

The specimens studied are deposited in the herbarium of the Institute of Applied Ecology, Chinese Academy of Sciences, Shenyang, China (IFP). The microscopic procedure follows He & Dai (2012). Sections were studied at magnifications up to ×1000 using a Nikon Eclipse 80i microscope and phase contrast illumination. Measurements were made from sections stained with CB. In presenting basidiospore size variation, the 5% of the measurements excluded from each end of the range are shown in parentheses. The meanings of abbreviations are as follows: CB = cotton blue, CB– = acyanophilous, IKI = Melzer's reagent, IKI– = negative in Melzer's reagent, KOH = 5% potassium hydroxide, L = mean basidiospore length (arithmetic average of all basidiospores), W = mean basidiospore width (arithmetic average of all basidiospores), Q = range of L/W ratios from individual specimens, and n = number of basidiospores measured from given number of specimens. Special color terms follow Petersen (1996).

Phylogenetic analysis

Total DNA was extracted from specimens using Phire® Plant Direct PCR Kit (Finnzymes, Finland). Primers ITS5 and ITS4 (White et al. 1990) were used to amplify ITS sequences following PCR procedure set forth by Stöger et al. (2006). The DNA was sequenced at Beijing Genomics Institute. The most similar sequences were downloaded from GenBank NCBI (http://www.ncbi.nlm.gov) using the BLAST tool. Sequences were aligned using ClustalX (Thomson et al. 1997) and manually edited to allow maximum alignment and minimize gaps. Maximum parsimony and Bayesian analysis were applied to the ITS dataset. All characters were weighted and gaps were treated as missing data. Maximum parsimony analysis were carried in PAUP* (version 4.0b10; Swofford 2002). Trees were inferred using the heuristic search option with TBR branch swapping and 1,000 random sequence additions. Max-trees were set to 5000 and no-increase, branches of zero length were collapsed, and all parsimonious trees were saved. Nodal support was tested with bootstrap of 1000 replicates using the heuristic search option (TBR and MULTREES options on) (Felsenstein 1985). Descriptive tree statistics tree length (TL), consistency index (CI), retention index (RI), rescaled consistency index (RC), and homoplasy index (HI) were calculated for all trees generated under different optimality criteria. Bayesian analysis was carried in MrBayes3.1.2 (Ronquist & Huelsenbeck 2003) and models of evolution were identified using MrMODELTEST2.3 (Posada & Crandall 1998, Nylander 2004). Four simultaneous Markov chains were run starting from random trees and keeping one tree every 100th generation until the average standard deviation of split frequencies was below 0.01. The value of burn-in was set to discard 25% of trees when calculating the posterior probabilities. Bayesian posterior probabilities were obtained from the 50% majority rule consensus of the trees kept as support for nodes.



Fig. 1. Strict consensus tree showing the phylogeny of *Melanoderma disciforme* and related species generated by Maximum Likelihood and Bayesian analysis based on ITS sequences. Parsimony bootstrap values >50% and Bayesian posterior probabilities >0.50 are shown on the branches.

Phylogenetic results

Two ITS sequences were newly generated in this study (GenBank accession numbers KM521268 and KM521269). The ITS dataset contains 34 sequences representing 24 species, among which *Cinereomyces lindbladii* (Berk.) Jülich was selected as outgroup. The alignment comprised 601 characters, of which 297 were constant, 50 were variable but parsimony-uninformative, and 254 were parsimony-informative. Maximum parsimony analysis yielded one parsimonious tree (CI = 0.505, RI = 0.669, RC = 0.338, HI = 0.495). Bayesian analysis ran 5 million generations and resulted in average standard deviation of split frequencies = 0.007737. The 50% majority consensus tree generated by the Bayesian analysis showed a similar topology with the strict consensus MP tree. The strict consensus tree is shown in Fig. 1. Both bootstrap support value (BP) and Bayesian posterior probabilities (BPP) are shown at the nodes. In the phylogenetic tree, two sequences of *M. disciforme* were grouped together with full support (100% BP, 1.00 BPP), and formed a monophyletic lineage with *M. microcarpum* fully supported (100% BP and 1.00 BPP).

Taxonomy

Melanoderma disciforme H.S. Yuan, **sp. nov.** Fig. 2

MycoBank MB 810703

Differs from *Melanoderma microcarpum* by its bigger pores, absence of encrusted cystidia, distinctly smaller cystidioles, and shorter basidiospores.

Type: China. Yunnan Prov., Xishuangbanna, Jinghong County, Nabanhe Nat. Res., on fallen angiosperm branch, 15.VIII.2005, Yuan 1675 (**holotype**, IFP; GenBank ITS, KM521269).

Etymology: the epithet "*disciforme*" refers to the disk-like basidiocarps.

Basidiocarps: Perennial, effuse-reflexed, growing as round pulvinate patches at first, then fusing laterally, margin narrowly reflexed as a pileus, back attached on substrates, coriaceous when fresh, woody hard upon drying, without odor or taste when fresh; pileus up to 3 mm broad, 5 cm long, and 3 mm thick at base. Pileal surface cream to buff when young, becoming black with a thin crust when aged, concentrically zonate and sulcate, glabrous; margin obtuse, cream to buff, ≤0.5 mm wide. Pore surface white to cream when fresh, buff when dry; pores round to angular, 6–7 per mm, dissepiments thick, entire; sterile margin distinct, ≤0.3 mm wide. Context cream to buff, hard corky, ≤1 mm thick. Tubes cream to buff, corky, distinctly stratified, about 1 mm long for each layer.

Hyphal structure: Hyphal system dimitic; generative hyphae clamped; skeletal hyphae dextrinoid and strongly cyanophilous; tissue unchanged in KOH.

Context: Generative hyphae scarce, hyaline, thin-walled, rarely branched, 1.5–2.5 μm in diam; skeletal hyphae dominant, hyaline, thick-walled to almost solid, frequently branched, interwoven, 1.5–3 μm in diam.

Tubes: Generative hyphae scarce, hyaline, thin-walled, rarely branched, 2.2–3 μm in diam; skeletal hyphae dominant, hyaline, thick-walled to almost solid, occasionally branched, interwoven, 1–3.5 μm in diam. Cystidia absent; cystidioles fusiform, hyaline, thin-walled, 11–14 × 3–4 μm. Basidia clavate to barrel, with four sterigmata and a basal clamp connection, 10–13 × 5–6 μm; basidioles similar in shape to basidia, but slightly smaller. Polygonal crystals frequently present in trama.

Basidiospores: Cylindrical, hyaline, thin-walled, smooth, IKI–, CB–, (4.8–)4.9–5.3(–5.5) × (1.9–)2–2.3(–2.5) μm, L = 5.05 μm, W = 2.17 μm, Q = 2.32–2.34 (n = 34/2).

Type of rot: Causes a white rot in angiosperms.

Additional specimens examined: ***Melanoderma disciforme***: **China**. **Yunnan Prov.**, Xishuangbanna, Jinghong County, Nabanhe Nat. Res., fallen angiosperm branch, 15.VIII.2005, Yuan 1643 (IFP; GenBank ITS: KM521268).

Melanoderma microcarpum: **China**. **Hunan Prov.**, Yizhang County, Mangshan Forest Park, on fallen angiosperm trunk, 25.VI.2007, Dai 8116 (BJFC, holotype).



Fig. 2. *Melanoderma disciforme* microscopic structures (Yuan 1675, holotype). a. Basidiospores; b. Basidia and basidioles; c. Cystidioles; d. Hyphae from trama; e. Hyphae from subiculum.

Discussion

Melanoderma disciforme represents the second species in the genus. It is characterized by perennial effuse-reflexed basidiocarps, narrow pileus with a thin black crust on the pileal surface, fusiform cystidioles, and cylindrical

basidiospores. The molecular phylogeny and morphological evidences support it as a new species.

Melanoderma disciforme shares most of the morphological features with *M. microcarpum*, the type species of the genus, which differs by smaller pores (7–9 per mm), apically encrusted cystidia, distinctly larger cystidioles, and longer basidiospores.

Species in *Perenniporia*, *Megasporoporiella* B.K. Cui et al., *Datronia* Donk, and *Dichomitus* D.A. Reid are morphologically similar to *M. disciforme* by having a dimitic hyphal system with cyanophilous skeletal hyphae. However, *Perenniporia* species have thick-walled basidiospores, *Megasporoporiella* species lack a crust on the surface of the pileus, and *Datronia* and *Dichomitus* species have non-dextrinoid skeletal hyphae (Núñez & Ryvarden 2001, Li & Cui 2013). Additionally, phylogenetic analyses based on ITS sequences indicate that species sampled from these genera are not genetically close to *M. disciforme*.

Acknowledgements

The authors would like to express their deep thanks to Drs. Juliano Marcon Baltazar (Universidade Estadual de Maringá, Brazil) and Shuanghui He (Beijing Forestry University, China) for serving as pre-submission reviewers. This research was financed by the National Natural Science Foundation of China (Project Nos. 31170022 & 31470148).

Literature cited

Cui BK, Zhao CL, Dai YC. 2011. *Melanoderma microcarpum* gen. et sp. nov. (*Basidiomycota*) from China. Mycotaxon 116: 295–302. http://dx.doi.org/10.5248/116.295

Dai YC. 2010. *Hymenochaetaceae* (*Basidiomycota*) in China. Fungal Divers. 45: 131–343. http://dx.doi.org/10.1007/s13225-010-0066-9

Dai YC. 2011. A revised checklist of corticioid and hydnoid fungi in China for 2010. Mycoscience 52: 69–79. http://dx.doi.org/10.1007/s10267-010-0068-1

Dai YC. 2012. Polypore diversity in China with an annotated checklist of Chinese polypores. Mycoscience 53: 49–80. http://dx.doi.org/10.1007/s10267-011-0134-3

Dai YC, Cui BK, Yuan HS, Li BD. 2007. Pathogenic wood-decaying fungi in China. Forest Pathology 37: 105–120. http://dx.doi.org/10.1111/j.1439-0329.2007.00485.x

Felsenstein J. 1985. Confidence intervals on phylogenetics: an approach using bootstrap. Evolution 39: 783–791. http://dx.doi.org/10.2307/2408678

He SH, Dai YC. 2012. Taxonomy and phylogeny of *Hymenochaete* and allied genera of *Hymenochaetaceae* (*Basidiomycota*) in China. Fungal Divers. 56: 77–93. http://dx.doi.org/10.1007/s13225-012-0174-9

Li HJ, Cui BK. 2013. Taxonomy and phylogeny of the genus *Megasporoporia* and its related genera. Mycologia 105: 368–83. http://dx.doi.org/10.3852/12-114

Ma J, Wang Y, Ma LG, Zhang YD, Castañeda-Ruíz RF, Zhang XG. 2011. Three new species of *Neosporidesmium* from Hainan, China. Mycological Progress 10: 157–162. http://dx.doi.org/10.1007/s11557-010-0685-2

Núñez M, Ryvarden L. 2001. East Asian polypores 2. *Polyporaceae* s. lato. Synop. Fungorum 14: 170–522.

Nylander JAA. 2004. MrModeltest v2. Program distributed by the author. Evolutionary Biology Centre, Uppsala University.

Petersen JH. 1996. Farvekort. The Danish Mycological Society's colour-chart. Foreningen til Svampekundskabens Fremme, Greve.

Posada D, Crandall KA. 1998. Modeltest: testing the model of DNA substitution. Bioinformatics 14: 817–818. http://dx.doi.org/10.1093/bioinformatics/14.9.817

Ronquist F, Huelsenbeck JP. 2003. MRBAYES 3: bayesian phylogenetic inference under mixed models. Bioinformatics 19: 1572–1574. http://dx.doi.org/10.1093/bioinformatics/btg180

Stöger A, Schaffer J, RuppitschW. 2006. A rapid and sensitive method for direct detection of *Erwinia amylovora* in symptomatic and asymptomatic plant tissues by polymerase chain reaction. J. Phytopathol. 154: 469–473.

Swofford DL. 2002. PAUP*. Phylogenetic analysis using parsimony (*and other methods). Version 4.0b10. Sinauer Associates, Sunderland, Massachusetts.

Thomson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. 1997. The Clustal_X windows interface: Flexible strategies for multiple sequence alignment aided by quality analysis tools. Nucleic Acids Res. 25: 4876–4882. http://dx.doi.org/10.1093/nar/25.24.4876

White TJ, Burns T, Lee S, Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. 315–322, in: MA Innis et al. (eds). PCR protocols, a guide to methods and applications. Academic, San Diego.
http://dx.doi.org/10.1016/B978-0-12-372180-8.50042-1

Zhang K, Ma J, Wang Y, Zhang XG. 2009. Three new species of *Piricaudiopsis* from southern China. Mycologia 101: 417–422. http://dx.doi.org/10.3852/09-006

Zhou LW, Dai YC. 2012. Recognizing ecological patterns of wood-decaying polypores on gymnosperm and angiosperm trees in northeast China. Fungal Ecology 5: 230–235. http://dx.doi.org/10.1016/j.funeco.2011.09.005